



# 53<sup>rd</sup> Annual Meeting of the EAAP Cairo, Egypt 4-7. September 2002



## AI IN PIG BREEDING IN ESTONIA

*A. Tänavots, T. Kaart\* & O. Saveli  
Institute of Animal Science, Estonian Agricultural University,  
Kreutzwaldi 1, Tartu 51014, Estonia*

### Abstract

The role of a male breeding animal is highly significant in livestock breeding, particularly in pig production due to very rapid turnover rate of pigs. Application of AI is increasing from year to year: in 1997 6% of the total number of pigs was inseminated artificially, whereas in 2001 the percentage was large (46.5%). Data of 6601 sows and 1015 boars with 10 411 litters, obtained from database of Animal Recording Centre in 1999...2001, was used to analyze heritability of litter size and effect of mating method on fertility traits. The following breed combinations were investigated: Estonian Landrace (EL), Estonian Large White (ELW), Hampshire (H), Pietrain (Pi), EL♂xELW♀, ELW♂xEL♀ and Pi♂xH♀. 9.80 piglets per litter were born by using AI, which was significantly lower (-0.44) than in case of natural mating. Significantly smaller litter size was observed in purebred EL (-0.39) and ELW (-0.62) breeds by using AI ( $P < 0.001$ ). H and Pi♂xH♀ combinations had larger litters at birth when AI was used. Natural mating showed superiority among parities, giving significantly larger litters from 1<sup>st</sup> to 6<sup>th</sup> parities. A rapid increase in application of AI shows that farmers have calculated advantages of AI and found that even in case of smaller litter size, they do not lose their profit, as they can use better genetic material.

### Introduction

The role of a male breeding animal is highly significant in livestock breeding, particularly in pig production due to very rapid turnover rate of pigs. Application of AI is increasing from year to year in Estonia: in 1997 6% of the total number of pigs was inseminated artificially, whereas in 2001 the percentage was larger (46.5%). Pioneer of the AI introduction was Kehtna AI Station, where 3068 sows were inseminated artificially in 1981. Today, there are four AI stations in Estonia, whereas the largest, Tartu AI station, produced about 26 000 sperm doses last year. Currently there are 40 boars in the Tartu AI Station (16 Estonian Large White, 11 Estonian Landrace, 9 Pietrain and 4 Pietrain x Hampshire crossbred boars). Average breeding value of the boars is 120 points, being higher (155 points) in Norwegian Landrace Fram 4398. Farmers could achieve nucleus farm breeding improvement also on their own farm, by using superior boar semen. Last year some Norwegian Landrace boar semen was imported and its offspring have a good body condition. More and more Pietrain boars are used to produce slaughter pigs. Farmers have a possibility to use also Pietrain x Hampshire crossbred boars (Rätsep, 2001).

Sows' fertility depends on many various factors (Clark & Leman, 1986), and in the previous studies the authors of the present paper (Tänavots, A. 1998<sup>a</sup>, Tänavots, A. 1998<sup>b</sup>, Tänavots *et al.*, 2001) found the influence of breed, parity, season and year on the fertility in purebred and crossbred pigs in Estonia. Factors influencing sow fertility also include mating type. Several studies have shown significant effect of this factor (Ral *et al.*, 1978; King *et al.*, 1998), however, Flowers & Alhusen (1992) did not find any difference.

The objective of this retrospective study was to investigate heritability of litter size and the effect of mating type [natural mating (NM) vs. artificial insemination (AI)] on fertility traits.

## Material and Methods

Data analysed comprised 6601 sows and 1015 boars with 10 411 litters from 39 farms throughout Estonia; obtained from database of Animal Recording Centre in 1999...2001. Completed dataset included breed, insemination method, farm, parity, season of birth and year of birth, which was collected by PC program DB-Planer. The following breed (litter) combinations were investigated: Estonian Landrace (EL), Estonian Large White (ELW), Hampshire (H), Piatrain (Pi), EL♂xELW♀, ELW♂xEL♀ and Pi♂xH♀. The testing year was divided into four parts: spring - March, April, May; summer - June, July, August, fall - September, October, November and winter - December, January, February.

General Linear Model (GLM) was used to analyse dataset by SAS software (SAS Inst. Inc., 1991).

$$Y_{ijkemnl} = \mu + T_i + M_j + K_k + S_e + A_m + P_n + e_{ijkemnl},$$

Y = dependent variable;

$\mu$  = general mean;

$T_i$  = farm (n=1...38);

$M_j$  = insemination method (n=1...2);

$K_k$  = boar (n=1...1015);

$S_e$  = birth season (n=1...4);

$A_m$  = birth year (n=1...3);

$P_n$  = breed (n=1...7);

$e_{ijkemnl}$  = random residual effect

The results are given as least-square means (Parring *et al.*, 1997). Level of significances expressed conventionally: \*\*\* - P<0.001, \*\* - P<0.01, \* - P<0.05, # - P<0.1.

## Results and Discussion

The average ratio of litters obtained through NM to litters obtained through AI was about 53%NM: 47%AI (figure 1), distribution of breed groups is shown in table 1.




Figure 1. Development of AI usage from 1999 to 2001

Table 1. Usage of AI in percentages, distributed by breeds

Traits	EL	ELW	H	Pi	ELxELW	ELWxEL	PixH
n	3428	4654	37	69	1515	682	26
AI	36.73	22.26	16.22	21.74	49.83	75.95	76.92
NM	63.27	77.74	83.78	78.26	50.17	24.05	23.08

From total 10 411 litters, 9.80 piglets per litter were born by using AI, which was significantly lower (-0.44) than in case of natural mating. Similar results were obtained by Tummaruk *et. al.* (2000). In their trials NM resulted in larger litters (0.2;  $P < 0.001$ ), compared with AI. By using AI a significantly smaller litter size was observed in purebred EL (-0.39) and ELW (-0.62) breeds ( $P < 0.001$ ) and little larger litters had H and Pi♂xH♀ combinations. As EL and ELW are the main breeds, used in Estonia, they have a large influence on total variation. In Swedish study, larger litters had also purebred Swedish Landrace and Swedish Yorkshire by using NM (Tummaruk *et. al.*, 2000). There is no explanation, why there are no differences in litter size between mating methods, while crossing white breeds.

Table 2. Differences between artificial insemination (AI) and natural mating (NM) among breeds

Traits	n	AI	NM	Difference	Significance
Total	10 411	9.80	10.25	-0.44	***
EL	3428	10.46	10.85	-0.39	***
ELW	4654	10.63	11.25	-0.62	***
H	37	10.19	9.12	1.07	n.s.
Pi	69	8.10	9.37	-1.27	n.s.
EL♂xELW♀	1515	10.52	10.64	-0.12	n.s.
ELW♂xEL♀	682	10.51	10.88	-0.38	n.s.
Pi♂xH♀	26	11.49	9.39	2.10	n.s.

As the effect of litter reduction, caused by AI, is considered in not significant on crossbreeding, it is suggested to use AI to produce slaughter pigs to achieve gain from better genetic material collected into AI stations.

Natural mating showed superiority among parities, giving significantly larger litters from 1<sup>st</sup> to 6<sup>th</sup> parities, whereas for the higher parities no significant difference was found. Compared with Tummaruk *et. al.* (2000) study, significantly smaller litter size was resulted for 1, 2 and 3 parities by using AI.

Table 3. Differences between artificial insemination (AI) and natural mating (NM) among parities

Traits	n	AI	NM	Difference	Significance
1	1774	9.01	9.40	-0.39	*
2	1928	9.83	10.09	-0.26	#
3	1940	10.31	10.56	-0.25	#
4	1607	10.71	11.12	-0.41	**
5	1240	10.42	11.18	-0.76	***
6	795	10.58	11.04	-0.45	*
7	489	10.86	11.01	-0.15	n.s.
8	323	11.06	11.30	-0.24	n.s.
9	174	10.89	11.29	-0.40	n.s.
10...15	141	10.58	10.96	-0.38	n.s.

The difference between NM and AI varied according to piglet breed and parity number.

A rapid increase in application of AI shows that farmers have calculated advantages of AI and found that even in case of smaller litter size, they do not lose their profit, as they can use better genetic material.

## References

- Clark, L. K., Leman, A.D. 1986. Factors that influence litter size in pigs. *Pig News Information*. 7, p. 303...310.
- Flower, W.L. & Alhusen, H.D. 1992. Reproductive performance and estimates of labor requirements associated with combinations of artificial insemination and natural service in swine. *Journal of Animal Science*. 70, p. 615...621.
- King, V.L., Koketsu, Y., Reeves, D., Xue, J.L. & Dial, G.D. 1998. Management factors associated with swine breeding-herd productivity in United States. *Prev. Vet. Med.* 35, p. 255...264.
- Parring, A-M., Vähi, M., Käärik, E. 1997. Statistilise andmetöötuse algõpetus. TÕ Mat. stat. inst. TÕ Kirjastus. p. 183...254.
- Rätsep, M. 2001. Kunstliku seemenduse kasutamine Eesti seafarmides kasvab järjekindlalt. *Tõuloomakasvatus* No. 4, p. 18...21.
- Ral, G., Andersson, K. & Kihlberg, M. 1978. Differences in litter size arising between artificially inseminated and naturally mated pigs in everyday production. *Swedish Journal of Agriculture, Res.* 8, p. 21...24.
- SAS. 1991. SAS User's guide: Statistics. SAS Inst. Inc., GARY, NC. 305 pp.
- Tummaruk, P., Lundeheim, N., Einarsson, S., Dalin, A.M. 2000. Reproductive performance of purebred Swedish Landrace and Swedish Yorkshire sows: II. effect of mating type, weaning-to-first-service interval and lactation length. *Acta Agriculturae Scandinavica. Section A, Animal Science*. 50:3, p. 217...224.
- Tänavots, A. 1998<sup>a</sup>. Crossbreeding influence on fertility traits of Estonian Large White sows. 4<sup>th</sup> Baltic Animal Breeding Conference, Tartu. p. 120...123.
- Tänavots, A. 1998<sup>b</sup>. Fertility and Lactation Yield of Large White Sows. Book of Abstracts of the 49<sup>th</sup> Annual Meeting of the EAAP. Warsaw, Poland, 24-27 August. p. 282.
- Tänavots, A., Saveli O., Kaart, T. 2001. Factors Affecting Meat Traits and Fertility of Pigs in Estonia. Book of Abstracts of the 52<sup>nd</sup> Annual Meeting of the EAAP, Budapest, Hungary, 26-29 August 2001. p. 300.

Acknowledgment: Estonian Science Foundation grant no. 0171417s99, Animal Recording Centre